



On seminormal monoid rings

Winfried Bruns^{a,*}, Ping Li^b, Tim Römer^a

^a *FB Mathematik/Informatik, Universität Osnabrück, 49069 Osnabrück, Germany*

^b *Department of Mathematics & Statistics, Queen's University, Kingston, ON K7L 3N6, Canada*

Received 6 June 2005

Available online 15 December 2005

Communicated by Steven Dale Cutkosky

Abstract

Given a seminormal affine monoid M we consider several monoid properties of M and their connections to ring properties of the associated affine monoid ring $K[M]$ over a field K . We characterize when $K[M]$ satisfies Serre's condition (S_2) and analyze the local cohomology of $K[M]$. As an application we present criteria which imply that $K[M]$ is Cohen–Macaulay and we give lower bounds for the depth of $K[M]$. Finally, the seminormality of an arbitrary affine monoid M is studied with characteristic p methods.

© 2005 Elsevier Inc. All rights reserved.

Contents

1. Introduction	362
2. Prerequisites	364
3. Seminormality and Serre's condition (S_2)	366
4. Local cohomology of monoid rings	369
5. The Cohen–Macaulay property and depth	377
6. Seminormality in characteristic p	381
7. Examples and counterexamples	382
References	386

* Corresponding author.

E-mail addresses: winfried@mathematik.uni-osnabrueck.de (W. Bruns), pingli@mast.queensu.ca (P. Li), troemer@mathematik.uni-osnabrueck.de (T. Römer).

1. Introduction

Let M be an affine monoid, i.e., M is a finitely generated commutative monoid which can be embedded into \mathbb{Z}^m for some $m \in \mathbb{N}$. Let K be a field and $K[M]$ be the affine monoid ring associated to M . Sometimes M is also called an affine semigroup and $K[M]$ a semigroup ring. The study of affine monoids and affine monoid rings has applications in many areas of mathematics. It establishes the combinatorial background for the theory of toric varieties, which is the strongest connection to algebraic geometry. In the last decades many authors have studied the relationship between ring properties of $K[M]$ and monoid properties of M . See Bruns and Herzog [3] for a detailed discussion and Bruns, Gubeladze and Trung [4] for a survey about open problems.

A remarkable result of Hochster [7] states that if M is normal, then $K[M]$ is Cohen–Macaulay. The converse is not true. It is a natural question to characterize the Cohen–Macaulay property of $K[M]$ for arbitrary affine monoids M in terms of combinatorial and topological information related to M . Goto, Suzuki and Watanabe [5] could answer this question for simplicial affine monoids. Later Trung and Hoa [18] generalized their result to arbitrary affine monoids. But the characterization is technical and not easy to check. Thus it is interesting to consider classes of monoids which are not necessarily simplicial, but nevertheless admit simple criteria for the Cohen–Macaulay property.

One of the main topics in the thesis [10] of the second author were seminormal affine monoids and their monoid rings. Recall that an affine monoid M is called *seminormal* if $z \in G(M)$, $2z \in M$ and $3z \in M$ imply that $z \in M$. Here $G(M)$ denotes the group generated by M . Hochster and Roberts [9, Proposition 5.32] noted that M is seminormal if and only if $K[M]$ is a seminormal ring. By a remarkable result of Gubeladze [6], finitely generated projective modules over $K[M]$ are free if M is seminormal. See Traverso [17] or Swan [16] for more details on seminormality. In general, there exist Cohen–Macaulay affine monoid rings which are not seminormal, and there exist seminormal affine monoid rings which are not Cohen–Macaulay. One of the main goals of this paper is to understand the problem in which cases $K[M]$ is Cohen–Macaulay for a seminormal affine monoid M . Another question is to characterize the seminormality property of affine monoids.

Let us go into more detail. Let R be a Noetherian ring and let N be a finitely generated R -module. The module N satisfies *Serre's condition* (S_k) if

$$\text{depth } N_{\mathfrak{p}} \geq \min\{k, \dim N_{\mathfrak{p}}\}$$

for all $\mathfrak{p} \in \text{Spec } R$. For trivial reasons Cohen–Macaulay rings satisfy Serre's condition (S_k) for all $k \geq 1$. The main result of [5] and a result of Schäfer and Schenzel [13] show that for a simplicial affine monoid M the ring $K[M]$ is Cohen–Macaulay if and only if $K[M]$ satisfies (S_2) . After some prerequisites we study in Section 3 the question to characterize the (S_2) property for $K[M]$ if M is a seminormal monoid. In the following let $C(M)$ be the cone generated by $M \subseteq \mathbb{Z}^m$. The main result in this section already appeared in the thesis of the second author [10] and states:

Theorem. *Let $M \subseteq \mathbb{Z}^m$ be a seminormal monoid and let F_1, \dots, F_t be the facets of $C(M)$. Then the following statements are equivalent:*

- (i) $K[M]$ satisfies (S_2) ;
- (ii) For all proper faces F of $C(M)$ one has

$$M \cap \text{int } F = \bigcap_{F \subseteq F_j} G(M \cap F_j) \cap \text{int } F;$$

- (iii) $G(M \cap F) = \bigcap_{F \subseteq F_j} G(M \cap F_j)$.

Here $\text{int } F$ denotes the relative interior of F with respect to the subspace topology on the affine hull of F . Let us assume for a moment that M is positive, i.e., 0 is the only invertible element in M . In order to decide whether $K[M]$ is a Cohen–Macaulay ring, one must understand the local cohomology modules $H_m^i(K[M])$ where m denotes the maximal ideal of $K[M]$ generated by all monomials X^a for $a \in M \setminus \{0\}$, because the vanishing and nonvanishing of these modules control the Cohen–Macaulayness of $K[M]$. Already Hochster and Roberts [9] noticed that certain components of $H_m^i(K[M])$ vanish for a seminormal monoid. Our result 4.3 in Section 4 generalizes their observation, and we can prove the following

Theorem. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine seminormal monoid such that $H_m^i(K[M])_a \neq 0$ for some $a \in G(M)$. Then $a \in -G(M \cap F) \cap F$ for a face F of $C(M)$ of dimension $\leq i$. In particular,*

$$H_m^i(K[M])_a = 0 \quad \text{if } a \notin -C(M).$$

As a consequence of this theorem and a careful analysis of the groups $H_m^i(K[M])$ we obtain in 4.7 that under the hypothesis of the previous theorem M is seminormal if and only if $H_m^i(K[M])_a = 0$ for all i and all $a \in G(M)$ such that $a \notin -C(M)$. Note that this result has a variant for the normalization, discussed in Remark 4.8.

Using further methods from commutative algebra we prove in Theorem 4.9:

Theorem. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid of rank d such that $K[M]$ satisfies (S_2) and $H_m^d(K[M])_a = 0$ for all $a \in G(M) \setminus (-C(M))$. Then M is seminormal.*

Since there are Cohen–Macaulay affine monoid rings which are not seminormal (like $K[t^2, t^3]$), one cannot omit the assumption about the vanishing of the graded components of $H_m^d(K[M])$ outside $-C(M)$.

In Section 5 we define the numbers

$$c_K(M) = \sup\{i \in \mathbb{Z}: K[M \cap F] \text{ is Cohen–Macaulay for all faces } F, \dim F \leq i\},$$

$$n(M) = \sup\{i \in \mathbb{Z}: M \cap F \text{ is normal for all faces } F, \dim F \leq i\}.$$

By Hochster’s theorem on normal monoids we have that $c_K(M) \geq n(M)$. The main result 5.3 in Section 5 are the inequalities

$$\text{depth } R \geq c_K(M) \geq \min\{n(M) + 1, d\}$$

for an affine seminormal monoid $M \subseteq \mathbb{Z}^m$ of rank d . Since seminormal monoids of rank 1 are normal, we immediately get that $\text{depth } K[M] \geq 2$ if $d \geq 2$. In particular, $K[M]$ is Cohen–Macaulay if $\text{rank } M = 2$.

One obtains a satisfactory result also for $\text{rank } M = 3$, which was already shown in [10] by different methods. In fact, in Corollary 5.6 we prove that $K[M]$ is Cohen–Macaulay for a positive affine seminormal monoid $M \subseteq \mathbb{Z}^m$ with $\text{rank } M \leq 3$ if $K[M]$ satisfies (S_2) . One could hope that $K[M]$ is always Cohen–Macaulay if M is seminormal and $K[M]$ satisfies (S_2) . But this is not the case and we present a counterexample in 7.1. The best possible result is given in 5.6: (S_2) is sufficient for $K[M]$ to be Cohen–Macaulay in the seminormal case if the cross-section of $C(M)$ is a simple polytope.

In Section 6 we study the seminormality of affine monoid rings with characteristic p methods. The main observation is that a positive affine monoid $M \subseteq \mathbb{Z}^m$ is seminormal if there exists a field K of characteristic p such that $K[M]$ is F -injective. This fact is a consequence of our analysis of the local cohomology groups of positive affine monoid rings. In 6.2 we give a precise description for which prime numbers p and fields of characteristic p , we have that $K[M]$ is F -injective. Implicitly, this result was already observed by Hochster and Roberts in [9, Theorem 5.33]. In fact, if $M \subseteq \mathbb{Z}^m$ is a positive affine seminormal monoid and K is a field of characteristic $p > 0$, then the following statements are equivalent:

- (i) The prime ideal (p) is not associated to the \mathbb{Z} -module $G(M) \cap \mathbb{R}F / G(M \cap F)$ for any face F of $C(M)$;
- (ii) R is F -split;
- (iii) R is F -pure;
- (iv) R is F -injective.

As a direct consequence we obtain that M is normal if the equivalent statements hold for every field K of characteristic $p > 0$.

In the last section we present examples and counterexamples related to the results of this paper. In particular, we will show that for every simplicial complex Δ there exists a seminormal affine monoid M such that the only obstruction to the Cohen–Macaulay property of $K[M]$ is exactly the simplicial homology of Δ . Choosing Δ as a triangulation of the real projective plane we obtain an example whose Cohen–Macaulay property depends on K . A similar result was proved by Trung and Hoa [18]. Our construction has the advantage of yielding a seminormal monoid M , and is geometrically very transparent.

2. Prerequisites

We recall some facts from convex geometry. Let X be a subset of \mathbb{R}^m . The *convex hull* $\text{conv}(X)$ of X is the set of convex combinations of elements of X . Similarly, the set $C(X)$ of positive linear combinations of elements of X is called the *cone* generated by X . By convention $C(\emptyset) = \{0\}$ and $\text{conv}(\emptyset) = \emptyset$. A cone C is called *positive* (or *pointed*) if 0 is the only invertible element in C . To an affine form α on \mathbb{R}^m (i.e., a polynomial of degree 1) we associate the affine hyperplane $H_\alpha = \alpha^{-1}(0)$, the closed half-space $H_\alpha^+ = \alpha^{-1}([0, \infty))$,

and the open half-space $H_\alpha^> = \alpha^{-1}(0, \infty)$. An intersection $P = \bigcap_{i=1}^n H_{\alpha_i}^+$ of finitely many closed half-spaces is called a *polyhedron*. A (proper) *face* of a polyhedron P is the (proper) intersection of P with a hyperplane H_β such that $P \subseteq H_\beta^+$. Also P is considered as a face of itself. A *facet* is a maximal proper face. Recall that there are only a finite number of faces. A *polytope* is a bounded polyhedron. The set $\text{conv}(F)$ is a polytope for every finite subset F of \mathbb{R}^m , and every polytope is of this form. A *cone* is a finite intersection of half-spaces of the form $H_{\alpha_i}^+$ where the α_i are linear forms (i.e., homogeneous polynomials of degree 1). The set $C(F)$ is a cone for every finite subset F of \mathbb{R}^m , and every cone is of this form. Let P be a polyhedron and F a face of P . Then we denote the relative interior of F with respect to the subspace topology on the affine hull of F by $\text{int } F$. Note that P decomposes into the disjoint union $\text{int } P$ of the (relative) interiors of its faces. For more details on convex geometry we refer to the books of Bruns and Gubeladze [2], Schrijver [14] and Ziegler [19].

An *affine monoid* M is a finitely generated commutative monoid which can be embedded into \mathbb{Z}^m for some $m \in \mathbb{N}$. We always use $+$ for the monoid operation. In the literature M is also called an *affine semigroup* in this situation. We call M *positive* if 0 is the only invertible element in M . Observe that M is positive if and only if $C(M)$ is pointed.

Let K be a field and $K[M]$ be the K -vector space with K -basis X^a , $a \in M$. The multiplication $X^a \cdot X^b = X^{a+b}$ for $a, b \in M$ induces a ring structure on $K[M]$ and this K -algebra is called the affine monoid ring (or algebra) associated to M . The embedding of M into \mathbb{Z}^m induces an embedding of $K[M]$ into the Laurent polynomial ring $K[\mathbb{Z}^m] = K[X_i^{\pm 1} : i = 1, \dots, m]$ where X_i corresponds to the i th element of the canonical basis of \mathbb{Z}^m . Note that $K[M]$ is a \mathbb{Z}^m -graded K -algebra with the property that $\dim_K K[M]_a \leq 1$ for all $a \in \mathbb{Z}^m$. It is easy to determine the \mathbb{Z}^m -graded prime ideals of $K[M]$. In fact every \mathbb{Z}^m -graded prime ideal is of the form $\mathfrak{p}_F = (X^a : a \in M, a \notin F)$ for a unique face F of $C(M)$ (see [3, Theorem 6.1.7] for a proof). In particular, the prime ideals of height 1 correspond to the facets of $C(M)$.

Recall that a Noetherian domain R is *normal* if it is integrally closed in its field of fractions. The normalization \bar{R} of R is the set of elements in the quotient field of R which are integral over R . An affine monoid M is called *normal*, if $z \in G(M)$ and $mz \in M$ for some $m \in \mathbb{N}$ imply $z \in M$. Here $G(M)$ is the group generated by M . It is easy to see that M is normal if and only if $M = G(M) \cap C(M)$. If M is an arbitrary submonoid of \mathbb{Z}^m , then its *normalization* is the monoid $\bar{M} = \{z \in G(M) : mz \in M \text{ for some } m \in \mathbb{N}\}$. By Gordan’s lemma \bar{M} is affine for an affine monoid M .

Hochster [7] proved that $K[M]$ is normal if and only if M is a normal monoid, in fact we have that $\bar{K[M]} = K[\bar{M}]$. In particular, Hochster showed that if M is normal, then $K[M]$ is a Cohen–Macaulay ring. One can characterize the Cohen–Macaulay property of $K[M]$ in terms of combinatorial and topological information associated to M . This amounts to an analysis of the \mathbb{Z}^m -graded structure of the local cohomology of $K[M]$; see Trung and Hoa [18] for a criterion of this type.

A Noetherian domain R is called *seminormal* if for an element x in the quotient field $Q(R)$ of R such that $x^2, x^3 \in R$ we have $x \in R$. The *seminormalization* ${}^+R$ of R is the intersection of all seminormal subrings S such that $R \subseteq S \subseteq Q(R)$. An affine monoid M is called *seminormal*, if $z \in G(M)$, $2z \in M$ and $3z \in M$ imply that $z \in M$. The seminormalization ${}^+M$ of M is the intersection of all seminormal monoids N such

that $M \subseteq N \subseteq G(M)$. It can be shown that ${}^+M$ is again an affine monoid. Hochster and Roberts [9, Proposition 5.32] proved that an affine monoid M is seminormal if and only if $K[M]$ is a seminormal ring. We frequently use the following characterization of seminormal monoids. See Reid and Roberts [11, Theorem 4.3] for a proof for positive monoids, but this proof works also for arbitrary affine monoids.

Theorem 2.1. *Let M be an affine monoid $M \subseteq \mathbb{Z}^m$. Then*

$${}^+M = \bigcup_{F \text{ face of } C(M)} G(M \cap F) \cap \text{int } F.$$

In particular, M is seminormal if and only if it equals the right-hand side of the equality.

3. Seminormality and Serre’s condition (S_2)

Let R be a Noetherian ring and let N be a finitely generated R -module. Recall that N satisfies *Serre’s condition* (S_k) if

$$\text{depth } N_{\mathfrak{p}} \geq \min\{k, \dim N_{\mathfrak{p}}\}$$

for all $\mathfrak{p} \in \text{Spec } R$. Affine monoid rings trivially satisfy (S_1) , since they are integral domains. We are interested in characterizing (S_2) for affine monoid rings.

While the validity of (S_k) in $K[M]$ may depend on the field K for $k > 2$, (S_2) can be characterized solely in terms of M , as was shown in [13].

Let F_1, \dots, F_t be the facets of $C(M)$ and let

$$M_i = \{a \in G(M): a + b \in M \text{ for some } b \in M \cap F_i\}$$

for $i = 1, \dots, t$. Note that the elements of M_i correspond to the monomials in the homogeneous localization $K[M]_{(\mathfrak{p}_{F_i})}$. We set

$$M' = \bigcap_{i=1}^t M_i.$$

Proposition 3.1. *Let M be an affine monoid $M \subseteq \mathbb{Z}^m$ and K a field. Then the following statements are equivalent:*

- (i) $K[M]$ satisfies (S_2) ;
- (ii) $M = M'$.

Observe that $(M')' = M'$. Thus $K[M']$ always satisfies (S_2) . For seminormal monoids M the equality $M = M'$ can be expressed in terms of the lattices $G(M \cap F)$ as we will see in Corollary 3.4. First we describe the monoids M_i under a slightly weaker condition.

Lemma 3.2. [10, Proposition 4.2.6] *Let $M \subseteq \mathbb{Z}^m$ be an affine monoid and let F_1, \dots, F_t be the facets of $C(M)$ with defining linear forms $\alpha_1, \dots, \alpha_t$. Then:*

- (i) $M_i \cap H_{\alpha_i} = G(M \cap F_i)$.
- (ii) If $G(M) \cap \text{int}C(M) \subseteq M$, then

$$M_i = (G(M) \cap H_{\alpha_i}^>) \cup G(M \cap F_i).$$

Proof. Every element in M_i is of the form $c = a - b$ for some $a \in M$ and $b \in M \cap F_i$. Hence $\alpha_i(c) \geq 0$ with equality if and only if $a \in M \cap F_i$. It follows that

$$M_i \subseteq (G(M) \cap H_{\alpha_i}^>) \cup G(M \cap F_i) \quad \text{and} \quad M_i \cap H_{\alpha_i} \subseteq G(M \cap F_i).$$

If $c \in G(M \cap F_i)$ and $c = a - b$ for some $a, b \in M \cap F_i$, then clearly by the definition of M_i we have that $c \in M_i$. Thus we see that $M_i \cap H_{\alpha_i} = G(M \cap F_i)$.

For (ii) it remains to show that if $c \in G(M) \cap H_{\alpha_i}^>$, then $c \in M_i$. Pick $d \in M \cap \text{int}F_i$ such that $\alpha_j(c + d) > 0$ for all $j \neq i$. Hence $c + d \in \text{int}C(M)$. But

$$c + d \in G(M) \cap \text{int}C(M) \subseteq M,$$

by the additional assumption in (ii). Thus $c \in M_i$. \square

In the following proposition we consider $C(M)$ as a face of itself.

Proposition 3.3. [10, Proposition 4.2.7] *Let M be an affine monoid $M \subseteq \mathbb{Z}^m$ and let F_1, \dots, F_t be the facets of $C(M)$. If $G(M) \cap \text{int}C(M) \subseteq M$, then*

$$M' = \bigcup_{F \text{ face of } C(M)} \left[\bigcap_{F \subseteq F_i} G(M \cap F_i) \cap \text{int}F \right]$$

with the convention that $\bigcap_{F \subseteq F_i} G(M \cap F_i) = G(M)$ if $F = C(M)$.

Proof. We apply 3.2 several times. By assumption we have

$$G(M) \cap \text{int}C(M) \subseteq M'.$$

Let F be a proper face of $C(M)$. Choose a facet F_j with defining linear form α_j . Either $\text{int}F \subseteq F_j$ and thus

$$\bigcap_{F \subseteq F_i} G(M \cap F_i) \cap \text{int}F \subseteq G(M \cap F_j) \cap \text{int}F \subseteq M_j,$$

or $\text{int}F$ is contained in $C(M) \cap H_{\alpha_j}^>$ and

$$\bigcap_{F \subseteq F_i} G(M \cap F_i) \cap \text{int}F \subseteq G(M) \cap H_{\alpha_j}^> \subseteq M_j.$$

Hence

$$\bigcap_{F \subseteq F_i} G(M \cap F_i) \cap \text{int } F \subseteq M'.$$

Note that

$$M' \cap \text{int } C(M) \subseteq G(M) \cap \text{int } C(M).$$

For a proper face F of $C(M)$ it follows from 3.2 that

$$M' \cap \text{int } F \subseteq \bigcap_{F \subseteq F_i} M_i \cap \text{int } F = \bigcap_{F \subseteq F_i} G(M \cap F_i) \cap \text{int } F.$$

All in all we see that

$$M' = \bigcup_{F \text{ face of } C(M)} \left[\bigcap_{F \subseteq F_i} G(M \cap F_i) \cap \text{int } F \right]. \quad \square$$

The equivalence of parts (ii) and (iii) in the following corollary was shown in [10, Theorem 4.2.14].

Corollary 3.4. *Let $M \subseteq \mathbb{Z}^m$ be an affine monoid, K a field, and let F_1, \dots, F_t be the facets of $C(M)$. If $G(M) \cap \text{int } C(M) \subseteq M$, then the following statements are equivalent:*

- (i) $K[M]$ satisfies (S_2) ;
- (ii) $M = M'$;
- (iii) For all proper faces F of $C(M)$ one has

$$M \cap \text{int } F = \bigcap_{F \subseteq F_j} G(M \cap F_j) \cap \text{int } F.$$

If M is seminormal, then the following is equivalent to (i)–(iii):

- (iv) $G(M \cap F) = \bigcap_{F \subseteq F_j} G(M \cap F_j)$.

Proof. The equivalence of (i) and (ii) was already stated in 3.1. The equivalence of (ii) and (iii) is an immediate consequence of 3.3.

In the seminormal case one has $G(F \cap M) \cap \text{int } F \subseteq M$ for all faces, so that (iv) implies (iii). For the converse implication one uses the fact that $G(M \cap F)$ is generated by its elements in $\text{int}(F)$ (see Bruns and Gubeladze [2]). \square

Remark 3.5. If M is seminormal, then we know from 2.1, that $G(M) \cap \text{int } C(M) \subseteq M$. Hence we can apply 3.2, 3.3 and 3.4 in this situation.

The corollary shows that a seminormal monoid satisfies (S_2) if and only if the restriction of the groups $G(M \cap F)$ happens only in the passage from $C(M)$ to its facets.

4. Local cohomology of monoid rings

For the rest of the paper K always denotes a field, and $M \subseteq \mathbb{Z}^m$ is an affine positive monoid of rank d . Recall that the seminormalization of M is

$${}^+M = \bigcup_{F \text{ face of } C(M)} G(M \cap F) \cap \text{int } F$$

and the normalization of M is

$$\bar{M} = G(M) \cap C(M).$$

In this section we want to compute the local cohomology of $K[M]$ and compare it with the local cohomology of $K[{}^+M]$ and $K[\bar{M}]$.

If M is a positive affine monoid, then $K[M]$ is a \mathbb{Z}^m -graded K -algebra with a unique graded maximal ideal \mathfrak{m} generated by all homogeneous elements of nonzero degree. By the local cohomology of $K[M]$ we always mean the local cohomology groups $H_{\mathfrak{m}}^i(K[M])$. Observe that ${}^+M$ and \bar{M} are also positive affine monoids. Since the K -algebras $K[{}^+M]$ and $K[\bar{M}]$ are finitely generated modules over $K[M]$ and the extensions of \mathfrak{m} are primary to their maximal ideals, the local cohomology groups of $K[{}^+M]$ and $K[\bar{M}]$ coincide with $H_{\mathfrak{m}}^i(K[{}^+M])$ and $H_{\mathfrak{m}}^i(K[\bar{M}])$, respectively. Because of this fact and to avoid cumbersome notation we always write $H_{\mathfrak{m}}^i(K[{}^+M])$ and $H_{\mathfrak{m}}^i(K[\bar{M}])$ for the local cohomology of $K[{}^+M]$ and $K[\bar{M}]$. The same applies to \mathbb{Z}^m -graded residue class rings of $K[M]$.

In the following R will always denote the ring $K[M]$, and thus ${}^+R$ and \bar{R} will stand for $K[{}^+M]$ and $K[\bar{M}]$, respectively.

Let F be a proper face of $C(M)$. Then $\mathfrak{p}_F = (X^a : a \in M, a \notin F)$ is a monomial prime ideal of R , and conversely, if \mathfrak{p} is a monomial prime ideal, then $F(\mathfrak{p}) = \mathbb{R}_+\{a \in M : X^a \notin \mathfrak{p}\}$ is a proper face of $C(M)$. These two assignments set up a bijective correspondence between the monomial prime ideals of R and the proper faces of $C(M)$.

Note that the natural embedding $K[M \cap F] \rightarrow K[M]$ is split by the face projection $K[M] \rightarrow K[M \cap F]$ that sends all monomials in F to themselves and all other monomials to 0. Its kernel is \mathfrak{p}_F . Therefore we have a natural isomorphism $K[M]/\mathfrak{p}_F \cong K[M \cap F]$.

The next lemma states a crucial fact for the analysis of the local cohomology of R . For this lemma and its proof we need the following notation. For $W \subseteq \mathbb{Z}^m$ we define

$$-W = \{-a : a \in W\}.$$

For a \mathbb{Z}^m -graded local Noetherian K -algebra R with $R_0 = K$ (like the monoid ring $K[M]$ for a positive affine monoid M) and a \mathbb{Z}^m -graded R -module N we set

$$N^\vee = \text{Hom}_K(N, K).$$

(Here we mean by $\text{Hom}_K(N, K)$ the homogeneous homomorphisms from N to K .) Note that N^\vee is again a \mathbb{Z}^m -graded R -module by setting

$$(N^\vee)_a = \text{Hom}_K(N_{-a}, K) \quad \text{for } a \in \mathbb{Z}^m.$$

Lemma 4.1. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid of rank d . The R -module $\bar{\omega}$ of R generated by the monomials X^b with $b \in \text{int } C(M) \cap G(M)$ is the canonical module of the normalization \bar{R} . If $a \in G(M)$, then*

$$H_m^i(\bar{\omega})_a \cong \begin{cases} 0 & \text{if } i < d \text{ or } a \notin -C(M), \\ K & \text{if } i = d \text{ and } a \in -C(M). \end{cases}$$

Proof. Danilov and Stanley showed that $\bar{\omega}$ is the canonical module of \bar{R} (see Bruns and Herzog [3, Theorem 6.3.5] or Stanley [15]). Thus $\bar{\omega}$ is Cohen–Macaulay of dimension d . This implies $H_m^i(\bar{\omega}) = 0$ for $i < d$. Furthermore, by graded local duality we have that

$$H_m^d(\bar{\omega})^\vee \cong \text{Hom}_{\bar{R}}(\bar{\omega}, \bar{\omega}) \cong \bar{R}$$

as \mathbb{Z}^m -graded modules. This concludes the proof. \square

For the central proofs in this paper it is useful to extend the correspondence between the faces of $C(M)$ and the monomial prime ideals of R to a bijection between the unions of faces of $C(M)$ and the monomial radical ideals. If \mathfrak{q} is a monomial radical ideal, then we let $F(\mathfrak{q})$ denote the union of the faces $F(\mathfrak{p})$ such that $\mathfrak{p} \supset \mathfrak{q}$, and if F is the union of faces, then the corresponding radical ideal \mathfrak{q}_F of R is just the intersection of all monomial prime ideals \mathfrak{p}_G such that $G \subseteq F$. We need the following lemma about monomial prime ideals of R .

Lemma 4.2. *Let $M \subseteq \mathbb{Z}^m$ be an affine monoid and F_1, \dots, F_t, G be faces of $C(M)$. Then:*

- (i) $\mathfrak{p}_{F_1 \cap \dots \cap F_t} = \mathfrak{p}_{F_1} + \dots + \mathfrak{p}_{F_t}$;
- (ii) $\mathfrak{p}_G + \bigcap_{i=1}^t \mathfrak{p}_{F_i} = \bigcap_{i=1}^t (\mathfrak{p}_G + \mathfrak{p}_{F_i})$.

Proof. Observe that the \mathfrak{p}_{F_i} are \mathbb{Z}^m -graded ideals of R , i.e., they are monomial ideals in this ring. In other words, their bases as K -vector spaces are subsets of the set of monomials X^a , $a \in M$. Using this fact it is easy to check the equalities claimed. \square

We are ready to prove a vanishing result for the local cohomology of seminormal monoid rings. Hochster and Roberts [9, Remark 5.34] already noticed that certain “positive” graded components of $H_m^i(R)$ vanish for a seminormal monoid. We can prove a much more precise statement.

Theorem 4.3. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine seminormal monoid and $R = K[M]$. If $H_m^i(R)_a \neq 0$ for $a \in \overline{G(M)}$, then $a \in -\overline{M} \cap \overline{F}$ for a face F of $C(M)$ of dimension $\leq i$. In particular,*

$$H_m^i(R)_a = 0 \quad \text{if } a \notin -C(M).$$

Proof. The assertion is trivial for $\text{rank } M = 0$. Thus assume that $d = \text{rank } M > 0$. Since M is seminormal, $\text{int } C(M) \cap G(M)$ is contained in M . Thus $\bar{\omega}$, which as a K -vector space is generated by the monomials X^a with $a \in \text{int } C(M) \cap G(M)$, is an ideal of R . Now consider the exact sequence

$$0 \rightarrow \bar{\omega} \rightarrow R \rightarrow R/\bar{\omega} \rightarrow 0.$$

By Lemma 4.1 the long exact local cohomology sequences splits into isomorphisms

$$H_m^i(R) \cong H_m^i(R/\bar{\omega}) \quad \text{for } i < d - 1 \tag{1}$$

and the exact sequence

$$0 \rightarrow H_m^{d-1}(R) \rightarrow H_m^{d-1}(R/\bar{\omega}) \rightarrow H_m^d(\bar{\omega}) \rightarrow H_m^d(R) \rightarrow 0. \tag{2}$$

The local cohomology of $\bar{\omega}$ has been determined in 4.1. Thus

$$H_m^d(R)_a = 0 \quad \text{for } a \notin -C(M).$$

This takes care of the top local cohomology. For the lower cohomologies we note that

$$R/\bar{\omega} \cong R / \bigcap_{i=1}^t \mathfrak{p}_{F_i}$$

where F_1, \dots, F_t are the facets of $C(M)$.

Therefore it is enough to prove the following statement which generalizes the theorem: *let \mathfrak{q} be a monomial radical ideal of R ; if $H_m^i(R/\mathfrak{q})_{-a} \neq 0$, then $a \in \overline{M} \cap \overline{G}$ for a face $G \subseteq F(\mathfrak{q})$ of dimension $\leq i$.*

The case $\mathfrak{q} = (0)$ has already been reduced to the case $\mathfrak{q} = \bar{\omega}$. So we can assume that $\mathfrak{q} \neq (0)$ and use induction on $\text{rank } M$ and on the number t of minimal monomial prime ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_t$ of \mathfrak{q} .

If $t = 1$, then $R/\mathfrak{p}_1 \cong K[M \cap F(\mathfrak{p}_1)]$. Now we can apply induction on $\text{rank } M$. Let $t > 1$. We set $\mathfrak{q}' = \bigcap_{j=1}^{t-1} \mathfrak{p}_j$. Then we have the standard exact sequence

$$0 \rightarrow R/\mathfrak{q} \rightarrow R/\mathfrak{q}' \oplus R/\mathfrak{p}_t \rightarrow R/(\mathfrak{q}' + \mathfrak{p}_t) \rightarrow 0.$$

The local cohomologies of R/\mathfrak{q}' and $R' = R/\mathfrak{p}_t$ are under control by induction. But this applies to $R/(\mathfrak{q}' + \mathfrak{p}_t)$, too. In fact, by Lemma 4.2 one has

$$R/(\mathfrak{q}' + \mathfrak{p}_t) \cong R / \bigcap_{j=1}^{t-1} (\mathfrak{p}_t + \mathfrak{p}_j) \cong R'/\mathfrak{q}''$$

where $\mathfrak{q}'' = \bigcap_{j=1}^{t-1} ((\mathfrak{p}_t + \mathfrak{p}_j)/\mathfrak{p}_t)$. Thus \mathfrak{q}'' is a monomial radical ideal of $R' \cong K[M \cap F(\mathfrak{p}_t)]$. Now it is enough to apply the long exact cohomology sequence

$$\dots \rightarrow H_m^{i-1}(R'/\mathfrak{q}'') \rightarrow H_m^i(R/\mathfrak{q}) \rightarrow H_m^i(R/\mathfrak{q}') \oplus H_m^i(R') \rightarrow \dots \quad \square$$

We describe a complex which computes the local cohomology of R . Writing R_F for the homogeneous localization $R_{(\mathfrak{p}_F)}$, let

$$L^\bullet(M) : 0 \rightarrow L^0(M) \rightarrow \dots \rightarrow L^t(M) \rightarrow \dots \rightarrow L^d(M) \rightarrow 0$$

be the complex with

$$L^t(M) = \bigoplus_{F \text{ face of } C(M), \dim F=t} R_F$$

and the differential $\partial : L^{t-1}(M) \rightarrow L^t(M)$ induced by

$$\partial_{G,F} : R_G \rightarrow R_F \text{ to be } \begin{cases} 0 & \text{if } G \not\subset F, \\ \epsilon(G, F) \cdot \text{nat} & \text{if } G \subset F, \end{cases}$$

where ϵ is a fixed incidence function on the face lattice of $C(M)$ in the sense of [3, Section 6.2]. In [3, Theorem 6.2.5] it was shown that $L^\bullet(M)$ is indeed a complex and that for an R -module N we have that

$$H_m^i(N) = H^i(L^\bullet(M) \otimes_R N) \quad \text{for all } i \geq 0.$$

Next we construct another, “smaller” complex which will be especially useful for the computation of the local cohomology of R if M is seminormal. Let

$${}^+L^\bullet(M) : 0 \rightarrow {}^+L^0(M) \rightarrow \dots \rightarrow {}^+L^t(M) \rightarrow \dots \rightarrow {}^+L^d(M) \rightarrow 0$$

be the complex with

$${}^+L^t(M) = \bigoplus_{F \text{ face of } C(M), \dim F=t} K[-\overline{M \cap F}]$$

and the differential ${}^+\partial : {}^+L^{t-1}(M) \rightarrow {}^+L^t(M)$ is induced by the same rule as ∂ above.

Proposition 4.4. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid, $a \in -C(M) \cap G(M)$. Then*

$$H_m^i(R)_a = H^i({}^+L(M))_a.$$

Proof. We know that $H_m^i(R)_a$ is the cohomology of the complex $L(M)_a$. To prove the claim it suffices to determine $(R_F)_a$ for a face F of $C(M)$ if $a \in -C(M) \cap G(M)$. It is an easy exercise to show that

$$(R_F)_a = K[-\overline{M \cap F}]_a,$$

where one has to use the fact that $\overline{M \cap F} = G(M \cap F) \cap F$. \square

It follows from 4.3 and 4.4 that the local cohomology of ${}^+R$ is a direct summand of the local cohomology of R as a K -vector space.

Corollary 4.5. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid. Then*

$$\bigoplus_{a \in -C(M) \cap G(M)} H_m^i(R)_a \cong \bigoplus_{a \in -C(M) \cap G(M)} H_m^i({}^+R)_a = H_m^i({}^+R).$$

Corollary 4.6. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid of rank d . Then:*

- (i) *If R is Cohen–Macaulay, then ${}^+R$ is Cohen–Macaulay.*
- (ii) *If $\text{depth } R \geq k$, then $\text{depth } {}^+R \geq k$.*
- (iii) *If R satisfies (S_k) , then ${}^+R$ satisfies (S_k) .*

Proof. It is well known that the Cohen–Macaulay property and depth can be read off the local cohomology groups. This is also true for Serre’s property (S_k) since we have that R satisfies (S_k) if and only if $\dim H_m^j(R)^\vee \leq j - k$ for $j = 0, \dots, \dim R - 1$ and an analogous characterization of Serre’s property (S_k) for ${}^+R$. (See Schenzel [12] for a proof of the latter fact.) \square

The results of this section allow us to give a cohomological characterization of seminormality for positive monoid rings.

Theorem 4.7. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid. Then the following statements are equivalent:*

- (i) *M is seminormal;*
- (ii) *$H_m^i(R)_a = 0$ for all i and all $a \in G(M)$ such that $a \notin -C(M)$.*

Proof. Consider the sequence

$$0 \rightarrow R \rightarrow {}^+R \rightarrow {}^+R/R \rightarrow 0$$

of finitely generated \mathbb{Z}^m -graded R -modules. Observe that $H_m^i(R)_a \cong H_m^i({}^+R)_a$ for $a \in -C(M)$. Thus it follows from the long exact cohomology sequence

$$\dots \rightarrow H_m^i(R) \rightarrow H_m^i({}^+R) \rightarrow H_m^i({}^+R/R) \rightarrow \dots$$

that $H_m^i(R)_a = 0$ for $a \notin -C(M)$ and all i if and only if $H_m^i({}^+R/R)_a = 0$ for all a and all i . This is equivalent to ${}^+R/R = 0$. Hence R and, thus, M are seminormal. \square

Remark 4.8. The previous results have variants for the normalization. If we restrict the direct sum in Corollary 4.5 to those a that belong to $-\text{int}C(M) \cap G(M)$ then the local cohomology of $K[{}^+M]$ must be replaced by that of \bar{R} . Moreover, the local cohomology of R vanishes in all degrees a outside $-\text{int}C(M)$ if and only if M is normal.

This follows by completely analogous arguments since we have that $H_m^i(\bar{R}) = 0$ for $i < d$ and $H_m^d(\bar{R})_a \neq 0$ if and only if $a \in -\text{int}C(M) \cap G(M)$.

With different methods than those used so far we can prove another seminormality criterion. It involves only the top local cohomology group, but needs a stronger hypothesis on M .

Theorem 4.9. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid of rank d . If R satisfies (S_2) and $H_m^d(R)_a = 0$ for all $a \in G(M) \setminus (-C(M))$, then M is seminormal.*

Proof. The assumption and 4.5 imply that the d th local cohomology of R and ${}^+R$ coincide as R -modules. Since M and therefore ${}^+M$ are positive, there exists a \mathbb{Z} -grading on R and ${}^+R$ such that both K -algebras are generated in positive degrees. We choose a common Noether normalization S of R and ${}^+R$ with respect to this \mathbb{Z} -grading.

Since R satisfies (S_2) it is a reflexive S -module. By 4.6 also ${}^+R$ satisfies (S_2) and is a reflexive S -module. In the following let ω_S be the canonical module of S which is in our situation just a shifted copy of S with respect to the \mathbb{Z} -grading. By graded local duality and reflexivity we get the following chain of isomorphisms of graded S -modules:

$$\begin{aligned} R &\cong \text{Hom}_S(\text{Hom}_S(R, S), S) \cong \text{Hom}_S(\text{Hom}_S(R, \omega_S), \omega_S) \\ &\cong \text{Hom}_S(H_m^d(R)^\vee, \omega_S) \cong \text{Hom}_S(H_m^d({}^+R)^\vee, \omega_S) \\ &\cong \text{Hom}_S(\text{Hom}_S({}^+R, \omega_S), \omega_S) \cong \text{Hom}_S(\text{Hom}_S({}^+R, S), S) \\ &\cong {}^+R. \end{aligned}$$

Hence $M = {}^+M$ is seminormal. \square

Again we can obtain a similar normality criterion if we replace $-C(M)$ by $-\text{int}C(M)$ in Theorem 4.9. In the rest of this section we further analyze the local cohomologies of R .

Proposition 4.10. *Let M be seminormal and $R = K[M]$. Then:*

- (i) $H_m^d(R)_{-a} \neq 0$ (and so of K -dimension 1) $\Leftrightarrow a \in \bar{M} \setminus \bigcup_F G(M \cap F)$ where F runs through the facets of $C(M)$.
- (ii) $H_m^{d-1}(R)_{-a} \neq 0 \Leftrightarrow a \in \partial C(M) \cap \bigcup_F G(M \cap F)$ and $\dim_K H_m^{d-1}(R/\bar{\omega})_{-a} \geq 2$ where F again runs through the facets of $C(M)$.
- (iii) R is Cohen–Macaulay $\Leftrightarrow R/\bar{\omega}$ is Cohen–Macaulay and $\dim_K H_m^{d-1}(R/\bar{\omega})_{-a} \leq 1$ for all $a \in G(M)$.

Proof. $H_m^d(R)$ is the cokernel of the map

$$\bigoplus_{F \text{ facet of } C(M)} K[-\overline{M \cap F}] \rightarrow K[-C(M) \cap G(M)],$$

which implies (i).

(ii) We have the exact sequence (2)

$$0 \rightarrow H_m^{d-1}(R) \rightarrow H_m^{d-1}(R/\bar{\omega}) \rightarrow H_m^d(\bar{\omega}) \xrightarrow{\pi} H_m^d(R) \rightarrow 0.$$

Lemma 4.1 and (i) yield the \mathbb{Z}^m -graded structure of $\text{Ker } \pi$: its nonzero graded components have dimension 1 and live in exactly the degrees $-a$ with $a \in C(M) \cap \bigcup_F G(M \cap F)$.

On the other hand, $H_m^{d-1}(R/\bar{\omega})$ can have nonzero components only in these degrees, as follows from the generalization of Theorem 4.3 stated in its proof. Thus $H_m^{d-1}(R)$ is limited to these degrees and is nonzero at $-a$ if and only if $\dim_K H_m^{d-1}(R/\bar{\omega})_{-a} \geq 2$.

(iii) The isomorphisms $H_m^i(R) \cong H_m^i(R/\bar{\omega})$ for $i < d - 1$ reduce the claim immediately to (ii). \square

Having computed the d th local cohomology of $K[M]$, we can easily describe the Gorenstein property of $K[M]$ in combinatorial terms:

Corollary 4.11. *Let M be seminormal and $R = K[M]$ Cohen–Macaulay. For each facet F of $C(M)$ let γ_F denote the index of the group extension $G(M \cap F) \subset G(M) \cap \mathbb{R}F$, and σ_F the unique \mathbb{Z} -linear form on $G(M)$ such that $\sigma_F(\bar{M}) = \mathbb{Z}_+$ and $\sigma_F(x) = 0$ for all $x \in F$. Then the following are equivalent:*

- (i) R is Gorenstein;
- (ii) (a) $\gamma_F \leq 2$ for all facets F of $C(M)$;
- (b) there exists $b \in \bar{M}$ such that

$$\begin{aligned} b \in F \setminus G(M \cap F), & \quad \text{if } \gamma_F = 2, \\ \sigma_F(b) = 1, & \quad \text{else.} \end{aligned}$$

Proof. The multigraded support of the K -dual ω_R of $H_m^d(R)$ is $N = \bar{M} \setminus \bigcup_F G(M \cap F)$, and R is Gorenstein if and only if there exists $b \in \bar{M}$ such that $N = b + M$, or, equivalently, $\omega_R = RX^b$. It remains to be shown that such b exists if and only if the conditions in (ii) are satisfied. We leave the exact verification to the reader. (Note that for each facet F there exists $c \in \bar{M} \cap \text{int}C(M) \subset M$ such that $\sigma_F(x) = 1$.) \square

Finally, we give an interpretation of $H_m^i(R)_{-a}$ for $a \in C(M)$ which will be useful in later sections.

Remark 4.12. Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid of rank d , $i \in \{0, \dots, d\}$, $a \in C(M) \cap G(M)$ and $\mathcal{F}(M, a) = \{F \text{ face of } C(M): a \in \overline{M \cap F}\}$. Then $H_m^i(R)_{-a}$ is the i th cohomology of the complex

$$C^\bullet(M, a) : 0 \rightarrow C^0(M, a) \rightarrow \dots \rightarrow C^t(M, a) \rightarrow \dots \rightarrow C^d(M, a) \rightarrow 0,$$

where

$$C^t(M, a) = \bigoplus_{G \in \mathcal{F}(M, a), \dim G=t} Ke_G$$

and the differential is given by $\partial(e_G) = \sum_{F \text{ face of } C(M), G \subseteq F} \varepsilon(G, F)e_F$. (Here ε is the incidence function on the face lattice of $C(M)$ which we fixed above to define the complex $L^\bullet(M)$.)

Theorem 4.13. Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid of rank d and $a \in C(M) \cap G(M)$. If the set $\mathcal{F}(M, a) = \{F \text{ face of } C(M) \mid a \in \overline{M \cap F}\}$ has a unique minimal element G , then

$$H_m^i(R)_{-a} = 0 \quad \text{for all } i = 0, \dots, d - 1.$$

Proof. It follows from 4.12 that $H_m^i(R)_{-a}$ is the i th cohomology of the complex

$$C^\bullet(M, a) : 0 \rightarrow C^0(M, a) \rightarrow \dots \rightarrow C^t(M, a) \rightarrow \dots \rightarrow C^d(M, a) \rightarrow 0,$$

where

$$C^t(M, a) = \bigoplus_{G \in \mathcal{F}(M, a), \dim G=t} Ke_G.$$

By taking a cross-section of $C(M)$ we see that the face lattice of $C(M)$ is the face lattice of a polytope (see [3, Proposition 6.1.8]). If $\mathcal{F}(M, a)$ has a unique minimal element, then this set is again the face lattice of a polytope P , as can be seen from Ziegler [19, Theorem 2.7]. Note that if $\mathcal{F}(M, a)$ has only one element, then P is the empty set. But this can only happen if $\mathcal{F}(M, a) = \{C(M)\}$ and then we have homology only in cohomological degree d . If $\mathcal{F}(M, a)$ has more than one element, then $C^\bullet(M, a)$ is the K -dual of a cellular resolution which computes the singular cohomology of P . A nonempty polytope is homeomorphic to a ball and thus the complex $C^d(M, a)$ is acyclic. Hence in this case $H_m^i(R)_{-a} = 0$ for $i = 0, \dots, d$, and this concludes the proof. \square

The following corollary collects two immediate consequences of Theorem 4.13.

Corollary 4.14. Let M be a positive affine monoid.

(i) Let F be the unique face of $C(M)$ such that $a \in \text{int } F$. If $a \in G(M \cap F)$, then

$$H_m^i(R)_{-a} = 0 \quad \text{for all } i = 0, \dots, d - 1.$$

(ii) Suppose that M is seminormal. Then R is Cohen–Macaulay if $\mathcal{F}(M, a)$ has a unique minimal element for all $a \in \bar{M}$.

Finally we note that nonzero lower local cohomologies must be large in the seminormal case.

Proposition 4.15. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine seminormal monoid. If $H_m^i(R)_a \neq 0$ for some $a \in -C(M)$, then $\dim_K H_m^i(R) = \infty$. In particular, if a seminormal monoid is Buchsbaum, then it must be Cohen–Macaulay.*

Proof. If $H_m^i(R)_a \neq 0$, then the complex $\mathcal{C}^*(M, a)$ has nontrivial cohomology in degree i . Consider the multiples ka for $k \in \mathbb{N}$. If $a \in \overline{M \cap F} = G(M \cap F) \cap F$, then $ka \in \overline{M \cap F}$ for all $k \in \mathbb{N}$. If $a \notin \overline{M \cap F}$, then there exist infinitely many k such that $ka \notin \overline{M \cap F}$. Since the face lattice of $C(M)$ is finite we can choose a sequence $(k_n)_{n \in \mathbb{N}}$ such that $k_n < k_{n+1}$ and $a \notin \overline{M \cap F}$ if and only if $k_n a \notin \overline{M \cap F}$. Thus $\mathcal{C}^*(M, a) = \mathcal{C}^i(M, k_n a)$ for all $n \geq 0$ which implies $H_m^i(R)_{k_n a} \neq 0$. Hence $\dim_K H_m^i(R) = \infty$.

If R is Buchsbaum, then $\dim_K H_m^i(R) < \infty$ for all $i < d$. Thus the local cohomology must vanish in this case for $i < d$ which implies that R is already Cohen–Macaulay. \square

5. The Cohen–Macaulay property and depth

If a seminormal monoid M fails to be normal by the smallest possible margin, then $K[M]$ is Cohen–Macaulay as the following result shows:

Proposition 5.1. *Let M be seminormal such that $M \cap F$ is normal for each facet F of $C(M)$. Then R is Cohen–Macaulay.*

Proof. It is enough to show that $\mathcal{F}(M, a)$ has a unique minimal element for all $a \in \bar{M}$. Let $a \in G(M) \cap C(M)$. If $a \notin \overline{M \cap G}$ for all facets G of $C(M)$, then $C(M)$ is the unique minimal element of $\mathcal{F}(M, a)$.

Otherwise we have $a \in \overline{M \cap G} = M \cap G \subset M$ for some facet G of $C(M)$. We choose the unique face F' of $C(M)$ with $a \in \text{int } F'$. It follows that $a \in F' \cap M$, and F' is the unique minimal element of $\mathcal{F}(M, a)$. \square

Remark 5.2. Another, albeit more complicated proof of the proposition can be given as follows. The main result of Brun, Bruns and Römer [1] implies for $R = K[M]$ that

- (i) $R/\bar{\omega}$ is Cohen–Macaulay,
- (ii) $H_m^d(R/\bar{\omega}) = \bigoplus_F H_m^{\dim F}(K[M \cap F])$ where F runs through the proper faces of $C(M)$,

provided all the rings $K[M \cap F]$ are Cohen–Macaulay. If they are even normal, then the local cohomology modules in (ii) do not “overlap” because the \mathbb{Z}^m -graded support of $H_m^{\dim F}(K[M \cap F])$ is restricted to $-\text{int } F$, and the relative interiors of faces are pairwise disjoint. Now we can conclude from Proposition 4.10 that R is Cohen–Macaulay.

In general, without normality of the facets the local cohomology modules in (ii) will overlap (see Example 7.1). This limits all attempts to prove stronger assertions about the Cohen–Macaulay property in the seminormal case.

Using the results and techniques of Section 4, we can give lower bounds for the depth of seminormal monoid rings. Let $M \subseteq \mathbb{Z}^m$ be an affine seminormal monoid. We define

$$c_K(M) = \sup\{i \in \mathbb{Z}: K[M \cap F] \text{ is Cohen–Macaulay for all faces } F, \dim F \leq i\},$$

$$n(M) = \sup\{i \in \mathbb{Z}: M \cap F \text{ is normal for all faces } F, \dim F \leq i\}.$$

Observe that if $M \cap F$ is normal for a face F of $C(M)$, then also $M \cap G$ is normal for all faces $G \subseteq F$ of $C(M)$. Hence it would be enough to consider all i -dimensional faces of $C(M)$ in the definition of $n(M)$. However, as we will see in Section 7, this is not true for the Cohen–Macaulay property.

Theorem 5.3. *Let $M \subseteq \mathbb{Z}^m$ be an affine seminormal monoid of rank d , and $R = K[M]$. Then*

$$\text{depth } R \geq c_K(M) \geq \min\{n(M) + 1, d\}.$$

Proof. The proof of the first inequality follows essentially the same idea as that of Theorem 4.3.

The assertion is trivial for $\text{rank } M = 0$. Thus assume that $d = \text{rank } M > 0$. There is nothing to prove if $c_K(M) = d$. So we can assume that $c_K(M) < d$. Since M is seminormal, we can again use the exact sequence

$$0 \rightarrow \bar{\omega} \rightarrow R \rightarrow R/\bar{\omega} \rightarrow 0.$$

Since $\text{depth } \bar{\omega} = d$ according to Lemma 4.1, it is enough to show that $\text{depth } R/\bar{\omega} \geq c_K(M)$.

Again we write $\bar{\omega} = \bigcap_{j=1}^t \mathfrak{p}_{F_j}$ where F_1, \dots, F_t are the facets of $C(M)$. However, contrary to Theorem 4.3, the bound does not hold for arbitrary residue class rings with respect to monomial radical ideals \mathfrak{q} , since the combinatorial structure of the set $F(\mathfrak{q})$ may contain obstructions.

Therefore we order the facets F_1, \dots, F_t in such a way that they form a shelling sequence for the face lattice of $C(M)$. Such a sequence exists by the Brugesser–Mani theorem (applied to a cross-section polytope of $C(M)$). See [19, Lecture 8]. The generalization of the first inequality of the theorem to be proved is the following: *let F_1, \dots, F_t be a shelling sequence for $C(M)$ and let $u \in \{1, \dots, t\}$, then $\text{depth } R/\mathfrak{q} \geq \min\{d - 1, c_K(M)\}$ for $\mathfrak{q} = \bigcap_{j=1}^u \mathfrak{p}_{F_j}$.*

If $u = 1$, then $R/\mathfrak{p}_1 \cong K[M \cap F(\mathfrak{p}_1)]$. Now we can apply induction on rank M . Let $u > 1$. We set $\mathfrak{q}' = \bigcap_{j=1}^{u-1} \mathfrak{p}_{F_j}$. Again we have the standard exact sequence

$$0 \rightarrow R/\mathfrak{q} \rightarrow R/\mathfrak{q}' \oplus R/\mathfrak{p}_{F_u} \rightarrow R/(\mathfrak{q}' + \mathfrak{p}_{F_u}) \rightarrow 0.$$

Therefore

$$\text{depth } R/\mathfrak{q} \geq \min\{1 + \text{depth } R/(\mathfrak{q}' + \mathfrak{p}_{F_u}), \text{depth } R/\mathfrak{p}_{F_u}, \text{depth } R/\mathfrak{q}'\}.$$

By induction on u we have $\text{depth } R/\mathfrak{p}_{F_u}, \text{depth } R/\mathfrak{q}' \geq \min\{d - 1, c_K(M)\}$.

Now the crucial point is that $F_u \cap \bigcup_{j=1}^{u-1} F_j = \bigcup_{j=1}^{u-1} F_u \cap F_j$ is the union of certain facets G_1, \dots, G_v of F_u that form the starting segment of a shelling sequence for F_u (by the very definition of a shelling). As in the proof of Theorem 4.3 we have

$$R/(\mathfrak{q}' + \mathfrak{p}_{F_u}) \cong R / \bigcap_{j=1}^{t-1} (\mathfrak{p}_{F_u} + \mathfrak{p}_{F_j}) \cong R'/\mathfrak{q}''$$

where $\mathfrak{q}'' = \bigcap_{j=1}^{t-1} ((\mathfrak{p}_{F_u} + \mathfrak{p}_{F_j})/\mathfrak{p}_{F_u})$. Therefore \mathfrak{q}'' is the radical ideal of $R' = K[M \cap F_1]$ corresponding to the union G_1, \dots, G_v . By induction we have

$$\text{depth } R'/\mathfrak{q}'' \geq \min\{d - 2, c(M \cap F_u)\} \geq \min\{d - 2, c_K(M)\},$$

and this completes the proof for the inequality $\text{depth } R \geq c_K(M)$.

By Hochster’s theorem the second inequality holds if M itself is normal. Suppose that $n(M) < d$ and let F be a face of dimension $n(M) + 1$. Then we must show that $K[M \cap F]$ is Cohen–Macaulay. Thus the second inequality reduces to the claim that R is Cohen–Macaulay if the intersections $F \cap M$ are normal for all facets F of $C(M)$ (and M is seminormal). This has been shown in Proposition 5.1. \square

There is a general lower bound for the depth of seminormal monoid rings of rank ≥ 2 . It follows from the proposition since seminormal monoids of rank 1 are normal.

Corollary 5.4. *Let $M \subseteq \mathbb{Z}^m$ be an affine seminormal monoid of rank $d \geq 2$. Then*

$$\text{depth } R \geq 2.$$

In particular, R is Cohen–Macaulay if $d = 2$.

One could hope that seminormality plus some additional assumptions on M already imply the Cohen–Macaulay property of R . But most time this is not the case as will be discussed in Example 7.1. However, we will now show that Serre’s condition (S_2) implies the Cohen–Macaulay property of R if $C(M)$ is a simple cone (to be explained below). More generally, we want to show that simple faces of $C(M)$ cannot contain an obstruction to the Cohen–Macaulay property in the presence of (S_2) .

Let F be a proper face of $C(M)$. We call the face F *simple* if the partially ordered set $\{G \text{ face of } C(M) : F \subseteq G\}$ is the face lattice of a simplex. Observe that by [19, Theorem 2.7] the latter set is always the face lattice of a polytope, because the face lattice of $C(M)$ is the face poset of a cross-section of $C(M)$. Let F be a simple face of $C(M)$. It is easy to see that every face G of $C(M)$ containing the simple face F is also simple.

We say that $C(M)$ is simple if a cross-section polytope of $C(M)$ is a simple polytope. This amounts to the simplicity of all the edges of $C(M)$. (Note that the apex $\{0\}$ is a simple face if and only if $C(M)$ is a simplicial cone.)

Proposition 5.5. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine seminormal monoid such that R satisfies (S_2) . Let $a \in G(M) \cap C(M)$ and $a \in \text{int } F$ for a proper face F of $C(M)$. If $H_m^i(R)_{-a} \neq 0$ for some i , $0 \leq i \leq d - 1$, then F is not a simple face of $C(M)$.*

Proof. Assume that F is a simple face. Consider the intersection

$$H = \bigcap_{G \text{ face of } C(M), F \subseteq G, a \in \overline{M \cap G}} G$$

which is a simple face containing F because F is simple. Let F_1, \dots, F_t be the facets of $C(M)$. For each facet F_j such that $H \subseteq F_j$ there exists a face G of $C(M)$ with $F \subseteq G$, $a \in \overline{M \cap G}$ such that $G \subseteq F_j$ because H is simple. This follows from the fact that the partially ordered set $\{L : L \text{ is a face of } C(M), H \subseteq L\}$ is the face poset of a simplex, and for a simplex the claim is trivially true. We observe that $a \in G(M \cap G) \subseteq G(M \cap F_j)$ for those facets F_j with $H \subseteq F_j$. By Corollary 3.4 we have

$$G(M \cap H) = \bigcap_{H \subseteq F_j} G(M \cap F_j).$$

Therefore $a \in G(M \cap H) \cap H = \overline{M \cap H}$.

All in all we get that the set $\mathcal{F}(M, a) = \{L \text{ face of } C(M) : a \in \overline{M \cap L}\}$ has the unique minimal element H , and 4.13 implies that

$$H_m^i(R)_{-a} = 0$$

which is a contradiction to our assumption. Thus F is not a simple face of $C(M)$. \square

The latter result gives a nice Cohen–Macaulay criterion in terms of $C(M)$ for a seminormal monoid. It implies Theorem 4.4.7 in [10], and can be viewed as a variant of the theorem by Goto, Suzuki and Watanabe [5] by which (S_2) implies the Cohen–Macaulay property of R if $C(M)$ is simplicial.

Corollary 5.6. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine seminormal monoid such that $R = K[M]$ satisfies (S_2) and such that $C(M)$ is a simple cone. Then R is Cohen–Macaulay for every field K . In particular, if $\text{rank } M \leq 3$, then R is Cohen–Macaulay.*

Proof. Every proper face of $C(M)$, with the potential exception of $\{0\}$, is simple. Thus it follows from 4.3 and 5.5 that $H_m^i(R)_{-a} = 0$ for $a \neq 0$ and $i = 0, \dots, d - 1$. For $a = 0$ this results from Corollary 4.14. Hence R is Cohen–Macaulay.

If $\text{rank } M \leq 3$, then the cross-section of $C(M)$ is a polytope of dimension ≤ 2 , which is always simple. Thus we can apply the corollary. \square

We will point out in Remark 7.2 that the corollary is the best possible result if one wants to conclude the Cohen–Macaulay property of R only from the seminormality of M and the validity of (S_2) .

6. Seminormality in characteristic p

In this section we study local cohomology properties of seminormal rings in characteristic $p > 0$. Let K be a field with $\text{char } K = p > 0$. In this situation we have the Frobenius homomorphism $F : R \rightarrow R, f \mapsto f^p$. Through this homomorphism R is an $F(R)$ -module. Now R is called F -injective if the induced map on the local cohomology $H_m^i(R)$ is injective for all i . It is called F -pure if the extension $F(R) \rightarrow R$ is pure, and F -split if $F(R)$ is a direct $F(R)$ -summand of R . In general we have the implications

$$F\text{-split} \Rightarrow F\text{-pure} \Rightarrow F\text{-injective}.$$

For example see [3] for general properties of these notions.

Proposition 6.1. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid. If there exists a field K of characteristic p such that R is F -injective, then M is seminormal.*

Proof. Assume that there exists an $a \in G(M)$, $a \notin -C(M)$, and an $i \in \{0, \dots, \text{rank } M\}$ such that $H_m^i(R)_a \neq 0$. Since R is F -injective, it follows that $H_m^i(R)_{p^m \cdot a} \neq 0$ for all $m \in \mathbb{N}$. Write $R = S/I_M$ as a \mathbb{Z}^m -graded quotient of a polynomial ring S . Then by graded local duality $H_m^i(R)^\vee \cong \text{Ext}_S^{n-i}(R, \omega_S)$ is a finitely generated \mathbb{Z}^m -graded R -module. This implies that $H_m^i(R)_{p^m \cdot a} = 0$ for $m \gg 0$, which is a contradiction. Thus $H_m^i(R)_a = 0$ for all $a \notin C(M)$. It follows from 4.7 that M is seminormal. \square

If M is seminormal there exist only finitely many prime numbers such that R is not F -injective. Moreover, we can characterize this prime numbers precisely.

Proposition 6.2. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine seminormal monoid and let K be a field of characteristic $p > 0$. Then the following statements are equivalent:*

- (i) *The prime ideal (p) is not associated to the \mathbb{Z} -module $G(M) \cap \mathbb{R}F/G(M \cap F)$ for any face F of $C(M)$;*
- (ii) *R is F -split;*
- (iii) *R is F -pure;*
- (iv) *R is F -injective.*

Proof. (i) \Rightarrow (ii) We show by a direct computation that $F(R) = K^p[pM]$ is a direct $K^p[pM]$ -summand of R . Since $K^p[pM]$ is a $K^p[pM]$ -summand of $K[pM]$, it is enough to show that $K[pM]$ is a direct $K[pM]$ -summand of R . The monoid M is the disjoint union of the residue classes modulo $pG(M)$. This induces a direct sum decomposition of R as a $K[pM]$ -module. We claim that pM is the intersection of M and $pG(M)$. This will show the remaining assertion.

To prove the claim we have only to show that an element w of the intersection of M and $pG(M)$ is an element of pM . Write $w = pz$ for some $z \in G(M)$. We have that $w \in \text{int } F$ for a face F of $C(M)$. Then p annihilates the element $z \in G(M) \cap \mathbb{R}F$ modulo $G(M \cap F)$. By assumption p is a nonzero-divisor on that module. Thus $z \in G(M \cap F)$. Since $z \in G(M \cap F) \cap \text{int } F$ and M is seminormal we have that $z \in M$ by 2.1. Hence $w \in pM$ as desired.

(ii) \Rightarrow (iii) and (iii) \Rightarrow (iv) This holds in general as was remarked above.

(iv) \Rightarrow (i) Assume that (p) is associated to some of the \mathbb{Z} -modules $G(M) \cap \mathbb{R}F / G(M \cap F)$ for the faces F of $C(M)$. Choose a maximal face F with this property. Choose an element $\bar{0} \neq \bar{a} \in G(M) \cap \mathbb{R}F / G(M \cap F)$ which is annihilated by p . We may assume that $a \in \text{int } F$.

It follows from 4.12 that $H_m^{\dim F+1}(R)_{-a} \neq 0$, since the poset $\mathcal{F}(M, a)$ consists all faces which contain F but not F itself. But $H_m^{\dim F+1}(R)_{-pa} = 0$, because $\mathcal{F}(pa)$ consists all faces which contain F including F itself. This poset is the face poset of a polytope and thus acyclic. We have derived a contradiction to the assumption that R is F -injective, because the map $H_m^{\dim F+1}(R)_{-a} \rightarrow H_m^{\dim F+1}(R)_{-pa}$ is not injective. \square

Corollary 6.3. *Let $M \subseteq \mathbb{Z}^m$ be a positive affine monoid. Then the following statements are equivalent:*

- (i) M is normal;
- (ii) R is F -split for every field K of characteristic $p > 0$;
- (iii) R is F -pure for every field K of characteristic $p > 0$;
- (iv) R is F -injective for every field K of characteristic $p > 0$.

Proof. It is easy to see that M is normal if and only if $G(M) \cap \mathbb{R}F = G(M \cap F)$ for all faces of $C(M)$. Thus (i) is equivalent to (ii), (iii) and (iv) by 6.1 and 6.2. \square

In a sense, it is inessential for the results of this section that K has characteristic p . In order to obtain variants that are valid for every field K , one has to replace the Frobenius endomorphism by the natural inclusion $K[pM] \rightarrow K[M]$.

7. Examples and counterexamples

In this section we present various examples and counterexamples related to the results of this paper. We choose a field K .

We saw in 5.6 that $R = K[M]$ is Cohen–Macaulay for a positive seminormal monoid M of rank $d \leq 3$. Since a Cohen–Macaulay ring always satisfies (S_2) one could hope that




Fig. 1.

seminormality and (S_2) already imply the Cohen–Macaulay property of R . This is not the case as the following example shows.

Example 7.1. For this example and the following one we fix some notation. Let P be a 3-dimensional pyramid with a square base embedded into \mathbb{R}^4 in degree 1. For example let P be the convex hull of the vertices given by

$$m_0 = (0, 0, 1, 1), \quad m_1 = (-1, 1, 0, 1), \quad m_2 = (-1, -1, 0, 1), \\ m_3 = (1, -1, 0, 1), \quad m_4 = (1, 1, 0, 1).$$

Figure 1 shows projections of the pyramid onto its base.

Let C be the cone generated by P , so that P is a cross-section of C . The facets of C are the cones

$$F_0 = C(m_1, m_2, m_3, m_4), \quad F_1 = C(m_0, m_1, m_2), \quad F_2 = C(m_0, m_2, m_3), \\ F_3 = C(m_0, m_3, m_4), \quad F_4 = C(m_0, m_1, m_4).$$

Let M be the monoid generated by all integer points of *even degree* in the facets F_1 and F_3 and all their faces, and all remaining integer points in the interior of all other faces of C including C . (The facets F_1 and F_3 have been shaded in the left diagram in Fig. 1.) Thus M is positive, $C(M) = C$ and M is not normal. It follows from 2.1 that M is seminormal and from 3.4 that R satisfies (S_2) .

We claim that R is not Cohen–Macaulay. Observe that all faces of C are simple except the face $C(m_0)$. Thus 4.3 and 5.5 imply that $H_m^i(R)_{-a}$ can be nonzero only for some $a \in C(m_0) \cap G(M)$. Choose an arbitrary a in the relative interior of $C(m_0) \cap G(M)$ of odd degree. The set $\mathcal{F}(M, a)$ introduced in 4.12 is

$$\{F_2, F_4, C\}$$

and we see that the complex $C^*(M, a)$ has cohomology in cohomological degree 3. Hence $H_m^3(R)_{-a} \neq 0$ and therefore R is not Cohen–Macaulay.

The reader may check that $R/\bar{\omega}$ is Cohen–Macaulay, but both $K[F_2 \cap M]$ and $K[F_4 \cap M]$ have nonzero third local cohomology in degree $-a$.

Remark 7.2. Example 7.1 can be generalized in the following way: if C is not a simple cone, then there exists a seminormal affine monoid M with $C = C(M)$ such that $K[M]$ satisfies (S_2) , but is not Cohen–Macaulay for any field K .

Next we consider the question whether the Cohen–Macaulay property or the (S_2) property are inherited by face projections. A counterexample to this claim is already given in [8, Example 2.2]. We can modify 7.1 a little bit to get the same result for seminormal monoids.

Example 7.3. With the same notation as in 7.1 let C be the cone over the pyramid P with facets F_0, \dots, F_4 . Now let M be the monoid generated by all integer points of *even degree* in the facet F_1 and all its faces (as indicated in the right diagram in Fig. 1), and all remaining integer points in the interior of all other faces of C . Thus M is positive, $C(M) = C$ and M is not normal. It still follows from 2.1 that M is seminormal and by 3.4 that R satisfies (S_2) . Since all proper faces of C except $C(m_0)$ are simple, we only have to check the vanishing of the local cohomology for points in the in $-\text{int}C(m_0)$. Let $a \in \text{int}C(m_0)$. If a has even degree, then

$$\mathcal{F}(M, a) = \{F \text{ face of } C: C(m_0) \subseteq F\}.$$

If a has odd degree, then

$$\mathcal{F}(M, a) = \{F_2 \cap F_3, F_3 \cap F_4, F_2, F_3, F_4, C\}.$$

In any case, we can check that the complex $\mathcal{C}^\bullet(M, a)$ is acyclic and therefore $H_m^i(R) = 0$ for $i < \text{rank } M$. Thus R is Cohen–Macaulay and must satisfy (S_2) .

But $K[M \cap F_3]$ has only depth 1, as can be seen from a similar discussion as for R . So it does not satisfy (S_2) . Hence neither the Cohen–Macaulay property, nor (S_2) are inherited by face projections of seminormal monoid rings.

Let Δ be a simplicial complex contained in the simplex Σ with vertex set V . We consider the dual simplex Σ^* whose facets correspond bijectively to the vertices $v \in V$ of Δ . Next we erect the pyramid Π over Σ^* with apex t . Then the faces of Π that contain t are in bijective correspondence with the faces G of Σ :

$$F \in \Sigma \iff F^* \in \Sigma^* \iff \tilde{F} = F^* \circ \{t\},$$

where \circ denotes the join. Observe that this correspondence reverses the partial order by inclusion. Choose a realization of Π as a rational polytope, also denoted by Π .

Next we plane off those faces of Π that correspond to the minimal nonfaces of Δ . For such a nonface G we choose a support hyperplane H of Π with $\Pi \cap H = \tilde{G}$. Moving this hyperplane by a sufficiently small rational displacement towards the interior of Π , and intersecting Π with the positive half-space of the displaced parallel hyperplane H' we obtain a polytope Π' such that exactly the faces F of Σ that do not contain G are preserved in Π' : $\tilde{F} \cap \Pi' \neq \emptyset \iff F \not\supseteq G$.

Repeating this construction for each minimal nonface of Δ we finally reach a polytope Π'' in which exactly the faces \tilde{F} , $F \in \Delta$, have survived in the sense that $F' = \tilde{F} \cap \Pi''$ is a nonempty face of Π'' .

Moreover, the only facets of Π'' containing F' are the facets $\{v\}'$ corresponding to the vertices $v \in F$. On the other hand, every face E of Π'' that is not of the form F' is contained in at least one “new” facet of Π'' created by the planing of Π .




Fig. 2. Planing off a face and the construction of Γ .

Note that Π is a simplex and therefore a simple polytope. The process by which we have created Π'' does not destroy simplicity if the displacements of the hyperplanes are sufficiently small and “generic.”

Set $d = \dim \Pi'' + 2$ and embed Π'' into $\mathbb{R}^{d-2} \times \{0\} \subset \mathbb{R}^{d-1}$. Then let Γ be the pyramid over Π'' with apex $v = (0, \dots, 0, 1)$. The construction of Γ that leads to the pyramid of Example 7.1 is illustrated in Fig. 2.

Note that all faces of Γ , except $\{v\}$, are simple. ($\{v\}$ is simple only if $\Delta = \Sigma$, or equivalently, $\Pi'' = \Pi$.)

In the last step we embed Γ into $\mathbb{R}^{d-1} \times \{1\} \subset \mathbb{R}^d$ by the assignment $x \mapsto (x, 1)$ and choose the cone $C = \mathbb{R}_+ \Gamma$. It has dimension d . The point v (in \mathbb{R}^d) has the coordinates $(0, \dots, 0, 1, 1)$. Therefore it has value 1 under the linear form $\text{deg}: \mathbb{R}^d \rightarrow \mathbb{R}$, $\text{deg}(y) = y_d$.

Set $L = \{a \in \mathbb{Z}^d: \text{deg}(a) \equiv 0 \pmod{2}\}$. To each facet F of C we assign the lattice

$$L_F = \begin{cases} \mathbb{R}F \cap \mathbb{Z}^d, & F = \mathbb{R}_+\{v\}' \text{ for some } v \in V, \\ \mathbb{R}F \cap L, & \text{else.} \end{cases}$$

Finally, we let M be the monoid formed by all $a \in C \cap \mathbb{Z}^d$ such that $a \in L_F$ for all facets F of C containing a . Clearly M is seminormal. Moreover, with the notation of Corollary 3.4, we have $M = M'$, since we have restricted the lattice facet-wise, and thus $K[M]$ satisfies (S_2) for all fields K .

Let $a \in \bar{M} = G(M) \cap C$. (By construction we have $G(M) = \mathbb{Z}^d$.) If the face F of C with $a \in \text{int}(F)$ is different from \mathbb{R}_+v , then it is a simple face of C , and $H_m^i(K[M])_{-a} = 0$ for all $i < n$ by Proposition 5.5. If $F = \mathbb{R}_+v$ and $\text{deg}(a) \equiv 0 \pmod{2}$, then we arrive at the same conclusion by Corollary 4.14. However, if $F = \mathbb{R}_+v$ and $\text{deg}(a) \equiv 1 \pmod{2}$, then the poset $\mathcal{F}(M, a)$ is isomorphic to the dual of Δ (as a poset). Hence the cochain complex $\mathcal{C}^\bullet(M, a)$ is isomorphic to the chain complex of Δ (up to shift). We have

$$H_m^i(K[M])_{-a} = \tilde{H}_{d-i-1}(\Delta; K), \quad i = 0, \dots, d. \tag{3}$$

Theorem 7.4. *Let Δ be a simplicial complex and K a field. Then there exists a seminormal monoid M of rank d with $M = M'$ and such that $R = K[M]$ has the following properties:*

- (i) For every $a \in \bar{M}$:
 - (a) $H_m^i(R)_{-a} = 0$ for all $i < d$; or
 - (b) $H_m^\bullet(R)_{-a}$ is given by (3).
- (ii) Moreover, case (b) holds true for at least one $a \in \bar{M}$.

(iii) *The following are equivalent:*

- (a) Δ is acyclic over K ;
- (b) R is Cohen–Macaulay.

If we choose Δ as a triangulation of the real projective plane, then we obtain a monoid algebra $K[M]$ which is Cohen–Macaulay if and only if $\text{char } K \neq 2$.

Acknowledgments

We are grateful to Aldo Conca for directing our attention toward the results of Hochster and Roberts in [9].

References

- [1] M. Brun, W. Bruns, T. Römer, Cohomology of partially ordered sets, math.AC/0502517.
- [2] W. Bruns, J. Gubeladze, Polytopes, rings, and K-theory, preprint.
- [3] W. Bruns, J. Herzog, Cohen–Macaulay Rings, revised ed., Cambridge Stud. Adv. Math., vol. 39, Cambridge Univ. Press, Cambridge, 1998.
- [4] W. Bruns, J. Gubeladze, N.V. Trung, Problems and algorithms for affine semigroups, Semigroup Forum 64 (2) (2002) 180–212.
- [5] S. Goto, N. Suzuki, K. Watanabe, On affine semigroup rings, Japan J. Math. (N.S.) 2 (1976) 1–12.
- [6] J. Gubeladze, Anderson’s conjecture and the maximal monoid class over which projective modules are free, Math. USSR-Sb. 63 (1) (1989) 165–180.
- [7] M. Hochster, Rings of invariants of tori, Cohen–Macaulay rings generated by monomials, and polytopes, Ann. of Math. (2) 96 (1972) 318–337.
- [8] M. Hochster, J.L. Roberts, Rings of invariants of reductive groups acting on regular rings are Cohen–Macaulay, Adv. Math. 13 (1974) 115–175.
- [9] M. Hochster, J.L. Roberts, The purity of the Frobenius and local cohomology, Adv. Math. 21 (1976) 117–172.
- [10] P. Li, Seminormality and the Cohen–Macaulay Property, PhD thesis, Kingston, 2004.
- [11] L. Reid, L.G. Roberts, Monomial subrings in arbitrary dimension, J. Algebra 236 (2) (2001) 703–730.
- [12] P. Schenzel, Zur lokalen Kohomologie des kanonischen Moduls, Math. Z. 165 (1979) 223–230.
- [13] U. Schäfer, P. Schenzel, Dualizing complexes of affine semigroup rings, Trans. Amer. Math. Soc. 322 (2) (1990) 561–582.
- [14] A. Schrijver, Theory of Linear and Integer Programming, Wiley, New York, 1998.
- [15] R.P. Stanley, Combinatorics and Commutative Algebra, second ed., Progr. Math., vol. 41, Birkhäuser, Basel, 1996.
- [16] R.G. Swan, On seminormality, J. Algebra 67 (1980) 210–229.
- [17] C. Traverso, Seminormality and Picard group, Ann. Scuola Norm. Sup. Pisa Sci. Fis. Mat. (3) 24 (1970) 585–595.
- [18] N.V. Trung, L.T. Hoa, Affine semigroups and Cohen–Macaulay rings generated by monomials, Trans. Amer. Math. Soc. 298 (1986) 145–167.
- [19] G.M. Ziegler, Lectures on Polytopes, Grad. Texts in Math., vol. 152, Springer-Verlag, Berlin, 1995.